

Figure 1

664760" 8/09/66




Figure 2

	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99																																																																																																																																																												
0	00000000	00000001	00000010	00000011	00000100	00000101	00000110	00000111	00001000	00001001	00001010	00001011	00001100	00001101	00001110	00001111	00010000	00010001	00010010	00010011	00010100	00010101	00010110	00010111	00011000	00011001	00011010	00011011	00011100	00011101	00011110	00011111	00100000	00100001	00100010	00100011	00100100	00100101	00100110	00100111	00101000	00101001	00101010	00101011	00101100	00101101	00101110	00101111	00110000	00110001	00110010	00110011	00110100	00110101	00110110	00110111	00111000	00111001	00111010	00111011	00111100	00111101	00111110	00111111	01000000	01000001	01000010	01000011	01000100	01000101	01000110	01000111	01001000	01001001	01001010	01001011	01001100	01001101	01001110	01001111	01010000	01010001	01010010	01010011	01010100	01010101	01010110	01010111	01011000	01011001	01011010	01011011	01011100	01011101	01011110	01011111	01100000	01100001	01100010	01100011	01100100	01100101	01100110	01100111	01101000	01101001	01101010	01101011	01101100	01101101	01101110	01101111	01110000	01110001	01110010	01110011	01110100	01110101	01110110	01110111	01111000	01111001	01111010	01111011	01111100	01111101	01111110	01111111	10000000	10000001	10000010	10000011	10000100	10000101	10000110	10000111	10001000	10001001	10001010	10001011	10001100	10001101	10001110	10001111	10010000	10010001	10010010	10010011	10010100	10010101	10010110	10010111	10011000	10011001	10011010	10011011	10011100	10011101	10011110	10011111	10100000	10100001	10100010	10100011	10100100	10100101	10100110	10100111	10101000	10101001	10101010	10101011	10101100	10101101	10101110	10101111	10110000	10110001	10110010	10110011	10110100	10110101	10110110	10110111	10111000	10111001	10111010	10111011	10111100	10111101	10111110	10111111	11000000	11000001	11000010	11000011	11000100	11000101	11000110	11000111	11001000	11001001	11001010	11001011	11001100	11001101	11001110	11001111	11010000	11010001	11010010	11010011	11010100	11010101	11010110	11010111	11011000	11011001	11011010	11011011	11011100	11011101	11011110	11011111	11100000	11100001	11100010	11100011	11100100	11100101	11100110	11100111	11101000	11101001	11101010	11101011	11101100	11101101	11101110	11101111	11110000	11110001	11110010	11110011	11110100	11110101	11110110	11110111	11111000	11111001	11111010	11111011	11111100	11111101	11111110	11111111


[illegible]

Figure 6




Figure 7




Figure 8




Figure 10




Figure 11

664760" 82096660




Figure 13

51

SHOULDER — 802c

Subscapularis

720

SYMPTOM PATTERNS — 804c

Local pain
Multiple nerves
Ulnar nerve
Bench press, push motions, swimming, popping shoulder, throwing, fall on arm, limited range with abduction and external rotation

FUNCTIONAL ANATOMY — 806c

Lesions at this site are often the result of heavy push motions. Competitive swimmers have this injury which leads to instability of the rotator cuff and impingement syndrome.

This muscle is essential for holding the head of the humerus posterior and inferior. With a lesion, the humerus translates anterior and superior causing popping and poor, inefficient biomechanics. This force couple is important to understand when treating shoulder injuries.

NOTE: When a lesion occurs it may affect the medial chord of the brachial plexus first. Then progress to all three chords of the brachial plexus when not treated appropriately. This is a common effect of the cumulative injury cycle.

Diagnosis — 808c

Myofascitis (rotator cuff syndrome, impingement syndrome)

Related Treatment Protocols — 810c

- 64 – Serratus anterior with subscapularis
- 71 – Brachial chords at the subscapularis
- 105 – Ulnar nerve at medial chord of brachial plexus and subscapularis

800c

TREATMENT — 812c

Contact — 814c

- Single thumb
- Single thumb with two finger backup

Pressure Tension — 816c

- High tension / low compression
- Delayed tension

Contact Motion — 818c

- Delayed internal slide

Patient Motion — 820c

- Passive first then active as soon as possible.

Instructions — 822c

1. Place the patient supine.
2. Grasp firmly and slightly abduct the humerus to a comfortable position for the patient that relieves tension on the subscapularis. Keep the patient's elbow bent 90 degrees or more. Do not externally rotate the humerus.
3. Palpate with your thumb past the latissimus dorsi and teres major to the anterior surface of the scapula. Feel the entire surface of the scapula that is covered by the subscapularis, move superior to the brachial chords, then medial into the axilla.
4. Contact inferior to the lesion and draw the skin and loose tissue to a point just supero-lateral to the area to be treated. Exert tension infero-medially along the muscle fiber orientation. Develop tension as the contact reaches the edge of the lesion.
5. Abduct the humerus until the arm overhead, keeping the elbow posterior as much as possible. Perform subsequent passes actively.

Notes — 824c

- Lack of gliding of the brachial chords will usually become evident during the second half of the motion.
- A defect in the fascia or a vein may feel like a thin, firm chord which arches across the body of the muscle. Do not attempt to break this down. Lighten the contact and pass over it to treat the remaining tissue during a pass.
- The muscle may be treated from the axilla to the medial border of the scapula where it meets the serratus anterior. Check for an adhesion between these two structures.

Figure 15




Figure 17

53

SHOULDER — 802d

Infraspinatus / Teres Minor with Deltoid

SYMPTOM PATTERNS — 804d

Local pain
Pulling / external rotation motions, difficulty reaching overhead and forward

FUNCTIONAL ANATOMY — 806d

During most motions of the shoulder, the infraspinatus and teres minor slide underneath the deltoid. Sometimes an adhesion develops at the lateral edge of the infraspinatus and teres minor, and the inferior edge of the posterior deltoid.

The infraspinatus and teres minor may also be adhered to the deltoid as it passes under it and around the humerus.

Diagnosis — 808d

Myofascitis

Related Treatment Protocols — 810d

- 45 – Deltoid
- 52 – Infraspinatus
- 54 – Teres minor

800d

TREATMENT — 812d

Contact — 814d

Single thumb

Pressure Tension — 816d

Delayed tension

Contact Motion — 818d

Internal and external slide

Patient Motion — 820d

Active

Instructions — 822d

1. Adduct the humerus with the patient prone.
2. Contact the inferior edge of the posterior deltoid and draw tension medially, arriving at the teres minor with your contact as much under the deltoid as possible.
3. Have the patient reach overhead, keeping the thumb pointing to the floor.

Infraspinatus Under Deltoid


1. Forward flex and externally rotate the humerus.
2. Contact the deltoid just anterior to the infraspinatus and traction the deltoid posterior along the fibers of the infraspinatus.
3. Maintain the traction while internally rotating the humerus.

Notes — 824d


- During patient motion, hold firm tension along the deltoid until it increases abruptly. Then draw your contact medially, trying to work between the muscles while letting the infraspinatus and teres slide underneath.

Figure 18


2000




Digital: 76




Long Thoracic: 79




Ulnar: 100, 101, 102, 103, 104, 105




Median: 80, 81, 82, 83, 84




Radial: 75, 92, 93, 94, 95, 96, 97, 98



Multiple: 71, 72, 88, 89, 90, 91



Musculocutaneous: 85, 86



Medial Antebrachial Cutaneous: 87


 Cutaneous Symptoms

Figure 20

71

SHOULDER — 802e

Brachial Chords at Subscapularis

2004

SYMPTOM PATTERNS — 804e

Ulnar nerve
Multiple nerves
Ulnar nerve / medial chord first, all three chords second, whole hand

FUNCTIONAL ANATOMY — 806e

The medial chord of the brachial plexus is usually the first of the chords to be entrapped with the subscapularis. If this process is allowed to worsen, the posterior and lateral chords follow. The brachial chords normally slide in excess of 1.5 centimeters during abduction of the humerus. When an adhesion occurs, it will exert a significant pull on the nerves during this motion.

The chords can be palpated by sliding your contact superior on the subscapularis to the roof of the axilla. The neurovascular sleeve should be palpable on the side of the contact thumb. It is possible to slide your contact to the anterior side of the sleeve and sometimes to the superior side as well. Take care to avoid applying undue compression on the nerves.

During treatment, it is common and desirable to duplicate the exact neurological symptoms. These usually subside in less than a minute, and the patient should be advised. As you palpate the nerve entrapment it is even possible to predict the symptom duplication.

Diagnoses — 808e

Peripheral nerve entrapment
Myofascitis

Related Treatment Protocols — 810e

51 – Subscapularis
90 – Neurovascular sleeve at coraco-pectoral tunnel
91 – Neurovascular sleeve at coracobrachialis

800e

TREATMENT — 812e

Contact — 814e

Single thumb

Pressure Tension — 816e

High tension / low compression

Contact Motion — 818e

Internal and external slide

Patient Motion — 820e

Active

Instructions — 822e


1. Place the patient supine and support the humerus with a firm comfortable grasp. Adduct the humerus to relieve tension on the subscapularis. Flex the elbow 90 degrees or more. Do not externally rotate the humerus. Palpate with your thumb past the latissimus dorsi and teres major to the anterior surface of the scapula, then superior to the brachial chords.
2. Contact inferior to the lesion. Draw the patient's skin and loose tissue to a point between the muscle and nerve. Exert tension along the nerve / muscle junction until sufficient tension is achieved.
3. Abduct the humerus, delaying motion of your contact until tension is felt to develop. Proceed within patient limits until the arm is raised overhead and the elbow is posterior as much as possible.

Notes — 824e


- Try to slide your contact between the muscle and nerve while holding the subscapularis medially and letting the nerve slide laterally. Make the second pass actively.
- Lack of brachial-chord sliding will usually become evident during the second half of the motion.

Figure 21


2300




Suprascapular: 99




Dorsal Scapular: 77, 78





Axillary: 74, 75





C-5 Nerve Root: 88, 89, Spine




C-8 Nerve Root: 88, 89, Spine



C-7 Nerve Root: 88, 89, Spine



C-6 Nerve Root: 88, 89, Spine



Internal Symptoms

Figure 23

664760" 84096260




Figure 24

83	<p>FOREARM — 802g</p> <p>Median Nerve at Pronator Teres</p>	800g
2008	<p>SYMPTOM PATTERNS — 804g</p> <p>Median nerve</p> <p>Local pain</p> <p>Symptoms of carpal tunnel syndrome, difficulty with grip, ache in forearm, edema</p>	804g
	<p>FUNCTIONAL ANATOMY — 806g</p> <p>The pronator teres is often injured with repetitive motion of the hands. It is the most common site of entrapment of the median nerve. Care must be taken to accurately work these tissues from one end to the other, including the ulnar and humeral head. After practice, it is possible to feel for relative motion between the median nerve and the pronator teres.</p> <p>Symptoms are usually duplicated as treatment reaches the entrapment site. At this site make every effort to ensure sliding of the nerve under the muscle. If symptoms of entrapment are severe, the release may have to be accomplished in steps. Work to patient tolerance.</p>	806g
	<p>Diagnoses — 808g</p> <p>Peripheral nerve entrapment</p> <p>Myofascitis</p>	808g
	<p>Related Treatment Protocols — 810g</p> <p>23 -- Flexor digitorum</p> <p>27 -- Pronator teres</p> <p>29 -- Lacertus fibrosis</p> <p>35 -- Brachioradialis</p> <p>82 -- Median nerve at flexor digitorum</p>	810g
	<p>TREATMENT — 812g</p> <p>Contact — 814g</p> <p>Single thumb</p> <p>Double thumb</p> <p>Pressure Tension — 816g</p> <p>High tension / low compression</p> <p>Contact Motion — 818g</p> <p>Internal and external slide</p> <p>Patient Motion — 820g</p> <p>Passive or active</p> <p>Instructions — 822g</p> <ol style="list-style-type: none"> 1. Pronate the patient's forearm and flex the elbow. 2. Draw tension on the pronator from the radial attachment proximally until moderate resistance is felt. 3. Have the patient slowly extend the elbow and supinate. As your contact nears the area where the muscle splits into a ulnar and humeral head, feel for the median nerve. Concentrate treatment in this area. <p>Notes — 824g</p> <ul style="list-style-type: none"> • Because this motion is difficult for the patient to learn, perform the first passes with the patient passive. Use active motions are subsequent to the patient learning the motion and a decrease in sensitivity in the severe cases. 	812g

Figure 26

90

SHOULDER / ARM / HAND — 802h

Neurovascular Sleeve at Coraco-Pectoral Tunnel

2004

SYMPTOM PATTERNS — 804h

Multiple nerves

Non nerve root symptoms, odd neurological patterns, all nerves, whole hand, difficult abduction/external rotation

FUNCTIONAL ANATOMY — 806h

The neurovascular sleeve holds the axillary artery, vein, and all the nerves into the upper extremity. This structure slides in excess of 1.5 centimeters through this thoracic outlet. When an adhesion occurs, it will exert a significant pull on the nerves during humeral abduction.

The sleeve may be palpated by sliding your contact superior on the subscapularis to the roof of the axilla. The neurovascular sleeve should be palpable on the side of the contact thumb. It is possible to slide your contact to the anterior side of the sleeve, and sometimes to the superior side of it as well. Avoid applying undue compression on the nerves.

Palpate along the posterior side of the pectoralis minor to the coracoid process at which point the sleeve is palpable. It is usually possible to work your contact between the pectoralis minor and the sleeve. The neurovascular sleeve may sometimes be traced and worked all the way to the clavicle and scalenes at the first rib.

Diagnoses — 808h

Peripheral nerve entrapment

Myofascitis

Related Treatment Protocols — 810h

44 – Coracobrachialis

47 – Pectoralis minor

51 – Subscapularis

85 – Neurovascular sleeve at the coracobrachialis

800h

TREATMENT — 812h

Contact — 814h

Single thumb

Pressure Tension — 816h

High tension / low compression

Contact Motion — 818h

Internal and external slide

Patient Motion — 820h

Active

Instructions — 822h

1. Orient the patient supine. Slightly abduct the humerus, supporting the patient's upper arm with a firm comfortable grasp. Flex the elbow 90 degrees or more, and do not externally rotate the humerus. Protract the shoulder.
2. Palpate with your thumb along the posterior side of the pectoralis minor to the sleeve and coracoid process. Exert medial tension along the nerve / muscle junction until sufficient tension is achieved.
3. Abduct the humerus, delaying supero-medial contact motion until tension develops. End with over 90 degrees abduction and maximum posterior extension. Stay within patient tolerance.

Notes — 824h

- Try to slide your contact between the muscle, or coracoid and nerve, while holding the muscle medially and letting the sleeve slide laterally. Follow the neurovascular sleeve as far toward the first rib and scalenes as possible.
- Lack of gliding of the sleeve will usually become evident during the second half of the motion.

Figure 28

09396078-04490




Figure 30

FOREARM / WRIST — 802j

Ulnar Nerve at Flexor Carpi Ulnaris

2022

SYMPTOM PATTERNS — 804j

Ulnar nerve
Local pain
Golfer's elbow, ulnar wrist pain, pain resting forearm

FUNCTIONAL ANATOMY — 806j

The ulnar nerve is not compromised in the cubital tunnel by compression. When the nerve is entrapped distal and proximal to the tunnel by the flexor carpi ulnaris and the medial intermuscular septum / triceps, it is stretched over the joint during elbow flexion. The nerve is flattened in the tunnel by the tension on either end.

When this happens, there is no decompression possible at the tunnel that will resolve the problem. It must be treated at the actual entrapment sites.

This muscle is important in ulnar nerve entrapment. It can, however, be injured without nerve involvement.

The area of concern is at the proximal end of the muscle where the nerve travels through it. In general, work the muscle first, then the interface between the nerve and the muscle on either side. Do not apply pressure directly on the nerve during treatment.

Because the flexor carpi ulnaris lies posterior to the elbow's axis of rotation, it lengthens during flexion. This is the opposite of the bulk of the forearm flexors.

Diagnoses — 808j

Peripheral nerve entrapment
Myofascitis

Related Treatment Protocols — 810j

22 – Pronator quadratus with flexor carpi ulnaris
42 – Medial intermuscular septum
71 – Brachial chords at subscapularis
72 – Brachial plexus at scalenes
90 – Neurovascular sleeve at coraco-pectoral tunnel
91 – Neurovascular sleeve at coracobrachialis
102 – Ulnar nerve at pronator quadratus and flexor carpi ulnaris

800j

TREATMENT — 812j

Contact — 814j

Multiple adjacent fingers
Double thumb

Pressure Tension — 816j

High tension / low compression

Contact Motion — 818j

Internal and external slide

Patient Motion — 820j

Active or passive

Instructions — 822j

1. Extend the patient's elbow and flex the wrist.
2. Contact on either side of the nerve at the proximal limit of the muscle, and draw tension from proximal to distal. Avoid excessive pressure on the ulnar nerve as it exits the cubital tunnel.
3. Flex the patient's elbow and extend the wrist as you draw your contact distally along the muscle/ nerve interface.

Figure 32